arXiv:math/0305340v1 [math.NT] 23 May 2003

Pair Correlation of the zeros of the Riemann zeta function in longer ranges

Tsz Ho Chan

February 1, 2008

Abstract

In this paper, we study a more general pair correlation function, $F_h(x,T)$, of the zeros of the Riemann zeta function. It provides information on the distribution of larger differences between the zeros.

1 Introduction

First of all, we assume the Riemann Hypothesis on the Riemann zeta function $\zeta(s)$ throughout this paper; $\rho = \frac{1}{2} + i\gamma$ denotes a non-trivial zero of the Riemann zeta function.

In the early 1970s, Hugh Montgomery considered the pair correlation function

$$F(x,T) = \sum_{0<\gamma,\gamma'\leq T} x^{i(\gamma-\gamma')} w(\gamma-\gamma') \text{ with } w(u) = \frac{4}{4+u^2}.$$

Here the sum is a double sum over the imaginary parts of the non-trivial zeros of $\zeta(s)$. He proved in [9] that, as $T \to \infty$,

$$F(x,T) \sim \frac{T}{2\pi} \log x + \frac{T}{2\pi x^2} \log^2 T$$

for $1 \leq x \leq T$ (actually he only proved for $1 \leq x \leq o(T)$ and the full range was done by Goldston [5]). He conjectured that

$$F(x,T) \sim \frac{T}{2\pi} \log T$$

for $T \leq x$ which is known as the Strong Pair Correlation Conjecture. From this, one has the (Weak) Pair Correlation Conjecture:

$$\sum_{\substack{0<\gamma,\gamma'\leq T \\ 0<\gamma-\gamma'\leq 2\pi\alpha/\log T}} 1 \sim \frac{T}{2\pi} \log T \int_0^\alpha 1 - \Big(\frac{\sin \pi u}{\pi u}\Big)^2 du.$$

which draws connections with random matrix theory.

The author studied these further in his thesis [1] (see also [2] and [3]) and derived more precise asymptotic formulas for $F(x,T)$ when x is in various ranges under the Twin Prime Conjecture **TPC** (see section 4). In the present paper, we generalize $F(x,T)$ further to

$$F_h(x,T) = \sum_{0<\gamma,\gamma'\le T} \cos\big((\gamma-\gamma'-h)\log x\big) w(\gamma-\gamma'-h).$$

Note that $F_h(x,T) = F_{-h}(x,T)$ and $F_0(x,T) = F(x,T)$. This leads to a better understanding of the distribution of larger differences between the zeros. Our main results are the following theorems: Here and throughout the paper, $\tilde{h} = |h|+1$.

Theorem 1.1. *For* $1 \le x \le \frac{T}{\log T}$,

$$\begin{aligned} F_h(x,T) =& \frac{T}{2\pi}\Big[\frac{4\cos(h\log x)}{4+h^2}\log x - \frac{8h\sin(h\log x)}{(4+h^2)^2}\Big] \\ &+ \frac{T}{2\pi x^2}\Big[\Big(\log\frac{T}{2\pi}\Big)^2 - 2\log\frac{T}{2\pi}\Big] + O(x\log x) + O\Big(\frac{\tilde{h}T}{x^{1/2-\epsilon}}\Big). \end{aligned}$$

Theorem 1.2. *Assume* **TPC**. *For* $M \ge 3$ *and* $\frac{T}{\log^M T} \le x$,

$$\begin{aligned} &F_h(x,T) \\ =&\frac{T}{\pi}\Big[\frac{2\cos(h\log x)}{4+h^2}\log x - \frac{4h\sin(h\log x)}{(4+h^2)^2}\Big] \\ +&\frac{T}{\pi}\int_1^\infty \Big[-\frac{2\cos(h\log x)}{4+h^2}\frac{1}{y} - \frac{4f(y)}{y^2}\cos(h\log x) + G_1(y) + G_2(y)\Big]\frac{\sin\frac{Ty}{x}}{\frac{Ty}{x}}dy \\ -&\frac{x}{\pi}\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{3\cos(h\log x)}{9+h^2} + \frac{h\sin(h\log x)}{9+h^2}\Big] \\ -&\frac{x}{\pi}\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{\cos(h\log x)}{1+h^2} - \frac{h\sin(h\log x)}{1+h^2}\Big] \\ +&\frac{T}{\pi}\sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2}\int_0^1 y\cos(h\log\frac{ky}{y})\frac{\sin\frac{Ty}{x}}{\frac{Ty}{x}}dy \\ +&O\Big(\tilde{h}\frac{x^{1+6\epsilon}}{T}\Big) + O(\tilde{h}x^{1/2+7\epsilon}) + O\Big(\tilde{h}\frac{x^2}{T^{2-2\epsilon}}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2}T}\Big). \end{aligned}$$

where $G_1(y)$ *and* $G_2(y)$ *are defined in Lemma 4.2.*

Theorem 1.3. *Assume* **TPC**. *For* $M \ge 3$ *and* $\frac{T}{\log^M T} \le x \le T$,

$$F_h(x,T) = \frac{T}{\pi}\Big[\frac{2\cos(h\log x)}{4+h^2}\log x - \frac{4h\sin(h\log x)}{(4+h^2)^2}\Big] + O(\tilde{h}x) + O\Big(\frac{\tilde{h}T}{\log^{M-2}T}\Big).$$

Theorem 1.4. *Assume* **TPC**. *For* $M \geq 3$ *and* $T \leq x \leq T^{2-29\epsilon}$,

$$F_h(x,T) = \frac{T}{2\pi} \log \frac{T}{2\pi e} \Big[\frac{4\cos{(h \log x)}}{4+h^2} \Big] + O\Big(\tilde{h} T \big(\frac{T}{x}\big)^{1/2-\epsilon} \Big) + O\Big(\frac{\tilde{h} T}{\log^{M-2} T} \Big).$$

For real α, let $F_h(\alpha) := (\frac{T}{2\pi} \log T)^{-1} F_h(T^\alpha, T)$. Then $F_h(\alpha) = F_h(-\alpha)$. Based on the above theorems, one may make the following

Conjecture 1.1. *For any arbitrary large* A, *as* $T \to \infty$,

$$F_h(\alpha) = \begin{cases} (1+o(1)) T^{-2\alpha} \log T + \alpha \frac{4\cos{(h \log T\alpha)}}{4+h^2} + o(1), & \text{if } 0 \leq \alpha \leq 1, \\ \frac{4\cos{(h \log T\alpha)}}{4+h^2} + o(1), & \text{if } 1 \leq \alpha \leq A. \end{cases}$$

By convolving $F_h(\alpha)$ with an appropriate kernel $\hat{r}(\alpha)$,

$$\Big(\frac{T}{2\pi} \log T \Big)^{-1} \sum_{0<\gamma,\gamma' \leq T} r\Big((\gamma - \gamma' - h) \frac{\log T}{2\pi} \Big) w(\gamma - \gamma' - h) = \int_{-\infty}^{+\infty} F_h(\alpha) \hat{r}(\alpha) \, d\alpha \tag{1}$$

where $\hat{r}(\alpha) = \int_{-\infty}^{\infty} r(u) e^{-2\pi i \alpha u} du$ for even $r(u)$ only. Conjecture 1.1 and (1) leads to

Conjecture 1.2. *For fixed* $\alpha > 0$,

$$\Big(\frac{T}{2\pi} \log T \Big)^{-1} \sum_{\substack{0<\gamma \neq \gamma' \leq T \\ |\gamma - \gamma' - h| \leq 2\pi\alpha / \log T}} 1 \sim \int_{-\alpha + h \log T/(2\pi)}^{\alpha + h \log T/(2\pi)} 1 - \frac{4}{4+h^2} \Big(\frac{\sin \pi u}{\pi u} \Big)^2 du.$$

Conjecture 1.3. *For* $0 < \alpha < \beta \ll \log T$,

$$\Big(\frac{T}{2\pi} \log T \Big)^{-1} \sum_{\substack{0<\gamma \neq \gamma' \leq T \\ 2\pi\alpha / \log T \leq \gamma - \gamma' \leq 2\pi\beta / \log T}} 1 \sim \int_{\alpha}^{\beta} 1 - \frac{1}{1 + (\pi u / \log T)^2} \Big(\frac{\sin \pi u}{\pi u} \Big)^2 du.$$

2 Some Lemmas

Lemma 2.1.

$$\begin{aligned} 2 \sum_{\gamma} \frac{x^{i(\gamma - t)}}{1 + (t-\gamma)^2} = & -\frac{1}{x} \sum_{n \leq x} \frac{\Lambda(n)}{n^{-1/2+it}} - x \sum_{n > x} \frac{\Lambda(n)}{n^{3/2+it}} + \frac{x^{1/2-it}}{\frac{1}{2} + it} + \frac{x^{1/2-it}}{\frac{3}{2} - it} \\ & + \frac{\log \tau}{x} + \frac{1}{x} \Big[\frac{\zeta'}{\zeta} \Big(\frac{3}{2} - it \Big) - \log 2\pi \Big] + O\Big(\frac{1}{x\tau} \Big) \end{aligned} \tag{2}$$

where the sum is over all the imaginary parts of the zeros of the Riemann zeta function, and $\tau = |t| + 2$. $\Lambda(n)$ *is von Mangoldt's lambda function.*

Proof: This is Lemma 2.2 in [2].

Write (2) as $\mathrm{Left}(x,t) = \mathrm{Right}(x,t)$. Let

$$\begin{aligned}
P(x,T) =& \frac{1}{x}\sum_{n\le x}\frac{\Lambda(n)}{n^{-1/2+it}} + x\sum_{n>x}\frac{\Lambda(n)}{n^{3/2+it}} - \frac{x^{1/2-it}}{\frac{1}{2}+it} - \frac{x^{1/2-it}}{\frac{3}{2}-it},\\
Q(x,T) =& \frac{\log\tau}{x},\\
R(x,T) =& \frac{1}{x}\Big[\frac{\zeta'}{\zeta}\Big(\frac{3}{2}-it\Big) - \log 2\pi\Big],\\
S(x,T) =& O\Big(\frac{1}{x\tau}\Big).
\end{aligned}$$

Lemma 2.2.

$$\begin{aligned}
&\int_0^T |\mathit{Left}(x,t) + \mathit{Left}(x,t-h)|^2 dt\\
=&2\pi F(x,T) + 2\pi F(x,T-h) + 4\pi F_h(x,T) + O(\log^3 T) + O(h\log^2 h).
\end{aligned}$$

Proof: This follows from page 188 of Montgomery [9] and the fact that $F(x,T) \ll T\log^2 T$.

Lemma 2.3. *For $x \ge 1$,*

$$\begin{aligned}
&\int_0^T |\mathit{Right}(x,t) + \mathit{Right}(x,t-h)|^2 dt\\
=&\int_0^T |P(x,t) + P(x,t-h)|^2 dt + \frac{4T}{x^2}\Big[\Big(\log\frac{T}{2\pi}\Big)^2 - 2\log\frac{T}{2\pi}\\
&+ \Big(\frac{1}{2}\sum_{n=1}^{\infty}\frac{\Lambda^2(n)(1+\cos(h\log n))}{n^3} + 2\Big)\Big] + O(\tilde{h}\log^2 T).
\end{aligned}$$

Proof: This is similar to the proof of Theorem 3.1 in [2].

Lemma 2.4. *For $x \ge 1$,*

$$\begin{aligned}
&4\pi F_h(x,T)\\
=&\int_0^T |P(x,t) + P(x,t-h)|^2 dt - \int_0^T |P(x,t)|^2 dt - \int_0^T |P(x,t-h)|^2 dt\\
&+ \frac{2T}{x^2}\Big[\Big(\log\frac{T}{2\pi}\Big)^2 - 2\log\frac{T}{2\pi} + \Big(\sum_{n=1}^{\infty}\frac{\Lambda^2(n)\cos(h\log n)}{n^3} + 2\Big)\Big] + O(\tilde{h}\log^3 T).
\end{aligned}$$

Proof: It follows from Lemma 2.2 and Lemma 2.3 as well as their special cases when $h = 0$.

Lemma 2.5. *For any sequence of complex numbers $\{a_n\}_{n=1}^{\infty}$ with $\sum_{n=1}^{\infty} n|a_n|^2 < \infty$,*

$$\int_0^T \Big|\sum_{n=1}^{\infty} a_n n^{-it}\Big|^2 dt = \sum_{n=1}^{\infty}|a_n|^2\big(T + O(n)\big).$$

Proof: This is Parseval's identity for Dirichlet series. See [10].

Lemma 2.6.

$$\begin{aligned}\sum_{n\le x}\Lambda^2(n)n =&\frac{1}{2}x^2\log x-\frac{1}{4}x^2+O(x^{1/2+\epsilon}).\\ \sum_{n>x}\frac{\Lambda^2(n)}{n^3} =&\frac{1}{2}\frac{\log x}{x^2}+\frac{1}{4x^2}+O\Big(\frac{1}{x^{5/2-\epsilon}}\Big).\end{aligned}$$

Proof: Use partial summation and the prime number theorem.

Lemma 2.7. *For any real a and b not both zero,*

$$\begin{aligned}\int e^{ax}\sin bx\,dx =&\frac{a}{a^2+b^2}e^{ax}\sin bx-\frac{b}{a^2+b^2}e^{ax}\cos bx.\\ \int e^{ax}\cos bx\,dx =&\frac{a}{a^2+b^2}e^{ax}\cos bx+\frac{b}{a^2+b^2}e^{ax}\sin bx.\\ \int xe^{ax}\sin bx\,dx =&\Big[\frac{ax}{a^2+b^2}-\frac{a^2-b^2}{(a^2+b^2)^2}\Big]e^{ax}\sin bx\\ &-\Big[\frac{bx}{a^2+b^2}-\frac{2ab}{(a^2+b^2)^2}\Big]e^{ax}\cos bx.\\ \int xe^{ax}\cos bx\,dx =&\Big[\frac{ax}{a^2+b^2}-\frac{a^2-b^2}{(a^2+b^2)^2}\Big]e^{ax}\cos bx\\ &+\Big[\frac{bx}{a^2+b^2}-\frac{2ab}{(a^2+b^2)^2}\Big]e^{ax}\sin bx.\end{aligned}$$

Proof: One can use $\int e^{(a+ib)x}dx$, $\int e^{(a-ib)}dx$, $\int xe^{(a+ib)x}dx$ and $\int xe^{(a-ib)x}dx$ which are simple to compute.

Lemma 2.8.

$$\begin{aligned}\frac{1}{x^2}\sum_{n\le x}\Lambda^2(n)n\cos(h\log n) =&\frac{2\cos(h\log x)}{4+h^2}\log x+\frac{h^2-4}{(4+h^2)^2}\cos(h\log x)\\ &+\frac{h\sin(h\log x)}{4+h^2}\log x-\frac{4h}{(4+h^2)^2}\sin(h\log x)\\ &+O\Big(\frac{\tilde{h}}{x^{1/2-\epsilon}}\Big).\\ x^2\sum_{n>x}\frac{\Lambda^2(n)}{n}\cos(h\log n) =&\frac{2\cos(h\log x)}{4+h^2}\log x-\frac{h^2-4}{(4+h^2)^2}\cos(h\log x)\\ &-\frac{h\sin(h\log x)}{4+h^2}\log x-\frac{4h}{(4+h^2)^2}\sin(h\log x)\\ &+O\Big(\frac{\tilde{h}}{x^{1/2-\epsilon}}\Big).\end{aligned}$$

Proof: We shall prove the first one. The other one is very similar. Let $A(x) = \frac{1}{x^2} \sum_{n \le x} \Lambda^2(n) n$. By integration by parts and Lemma 2.6,

$$\begin{aligned}
&\frac{1}{x^2} \sum_{n \le x} \Lambda^2(n) n \cos(h \log n) \\
=&\frac{A(x)}{x^2} \cos(h \log x) + \frac{h}{x^2} \int_1^x A(u) \frac{\sin(h \log u)}{u}\, du \\
=&\Big[\frac{1}{2} \log x - \frac{1}{4}\Big] \cos(h \log x) + \frac{h}{x^2} \int_1^x \Big[\frac{1}{2} \log u - \frac{1}{4}\Big] u \sin(h \log u)\, du \\
&+ O\Big(\frac{\tilde{h}}{x^{1/2-\epsilon}}\Big) \\
=&\Big[\frac{1}{2} \log x - \frac{1}{4}\Big] \cos(h \log x) + \frac{h}{x^2} \Big[\frac{1}{2} \int_0^{\log x} v e^{2v} \sin hv\, dv \\
&- \frac{1}{4} \int_0^{\log x} e^{2v} \sin hv\, dv\Big] + O\Big(\frac{\tilde{h}}{x^{1/2-\epsilon}}\Big)
\end{aligned}$$

which gives the desired result after applying Lemma 2.7 with $a = 2$ and $b = h$, and some algebra.

3 Proof of Theorem 1.1

First, note that

$$P(x,t) = \frac{1}{x^{1/2}} \Big[\sum_{n \le x} \Lambda(n) \Big(\frac{x}{n}\Big)^{-1/2+it} + \sum_{n > x} \Lambda(n) \Big(\frac{x}{n}\Big)^{3/2+it}\Big] + O\Big(\frac{x^{1/2}}{\tau}\Big).$$

Thus,

$$\begin{aligned}
P(x,t) + P(x,t-h) =&\frac{1}{x^{1/2}} \Big[\sum_{n \le x} \Lambda(n)(1 + n^{ih}) \Big(\frac{x}{n}\Big)^{-1/2+it} \\
&+ \sum_{n > x} \Lambda(n)(1 + n^{ih}) \Big(\frac{x}{n}\Big)^{3/2+it}\Big] + O\Big(\frac{x^{1/2}}{\tau}\Big)
\end{aligned}$$

So, the first integral in Lemma 2.4

$$\begin{aligned}
=&\frac{1}{x}\int_0^T \Big|\sum_{n\le x}\Lambda(n)(1+n^{ih})\Big(\frac{x}{n}\Big)^{-1/2+it}+\sum_{n>x}\Lambda(n)(1+n^{ih})\Big(\frac{x}{n}\Big)^{3/2+it}\Big|^2 dt\\
&+O\Big(\Big[\sum_{n\le x}\Lambda(n)\Big(\frac{x}{n}\Big)^{-1/2}+\sum_{n>x}\Lambda(n)\Big(\frac{x}{n}\Big)^{3/2}\Big]\int_0^T\frac{1}{\tau^2}dt\Big)+O\Big(\int_0^T\frac{x}{\tau^4}dt\Big)\\
=&\frac{1}{x}\sum_{n\le x}\Lambda^2(n)|1+n^{ih}|^2\Big(\frac{x}{n}\Big)^{-1}(T+O(n))\\
&+\frac{1}{x}\sum_{n>x}\Lambda^2(n)|1+n^{ih}|^2\Big(\frac{x}{n}\Big)^{3}(T+O(n))+O(x)\\
=&\frac{2T}{x^2}\sum_{n\le x}\Lambda^2(n)n\big(1+\cos{(h\log n)}\big)+2Tx^2\sum_{n>x}\frac{\Lambda^2(n)}{n^3}\big(1+\cos{(h\log n)}\big)\\
&+O(x\log x)
\end{aligned}$$

Similarly (or by setting $h=0$), each of the second and third integral in Lemma 2.4

$$=\frac{T}{x^2}\sum_{n\le x}\Lambda^2(n)n+Tx^2\sum_{n>x}\frac{\Lambda^2(n)}{n^3}+O(x\log x)$$

Therefore,

$$\begin{aligned}
4\pi F_h(x,T)=&2T\Big[\frac{1}{x^2}\sum_{n\le x}\Lambda^2(n)n\cos{(h\log n)}+x^2\sum_{n>x}\frac{\Lambda^2(n)}{n^3}\cos{(h\log n)}\Big]\\
&+\frac{2T}{x^2}\Big[\Big(\log\frac{T}{2\pi}\Big)^2-2\log\frac{T}{2\pi}\Big]\\
&+O\Big(\frac{T}{x^2}\Big)+O(\tilde{h}\log^3 T)+O(x\log x)\\
=&2T\Big[\frac{4\cos{(h\log x)}}{4+h^2}\log x-\frac{8h\sin{(h\log x)}}{(4+h^2)^2}\Big]\\
&+\frac{2T}{x^2}\Big[\Big(\log\frac{T}{2\pi}\Big)^2-2\log\frac{T}{2\pi}\Big]+O(x\log x)+O\Big(\frac{\tilde{h}T}{x^{1/2-\epsilon}}\Big)
\end{aligned}$$

by Lemma 2.8. The theorem follows after dividing through by 4π.

4 Twin Prime Conjecture and smooth weight

We shall use a quantitative form of the Twin Prime Conjecture **TPC** as follow: For any $\epsilon>0$,

$$\sum_{n=1}^N\Lambda(n)\Lambda(n+d)=\mathfrak{S}(d)N+O(N^{1/2+\epsilon})\text{ uniformly in }|d|\le N.$$

$\mathfrak{S}(d) = 2\prod_{p>2}\big(1-\frac{1}{(p-1)^2}\big)\prod_{p|d,p>2}\frac{p-1}{p-2}$ if d is even, and $\mathfrak{S}(d)=0$ if d is odd.

Let K and M be some large positive integers (K may depend on ϵ). Set $U=\log^M T$ and $\Delta = 1/(2^K U)$. We recall the smooth weight $\Psi_U(t)$ in [3] with:

1. support in $[-1/U, 1+1/U]$,
2. $0 \le \Psi_U(t) \le 1$,
3. $\Psi_U(t) = 1$ for $1/U \le t \le 1-1/U$,
4. $\Psi_U^{(j)}(t) \ll U^j$ for $j=1,2,...,K$.

This weight function satisfies the requirements in Goldston and Gonek [6]. One more thing to note is that

$$Re\hat{\Psi}_U(y) = \frac{\sin 2\pi y}{2\pi y}\Big(\frac{\sin 2\pi\Delta y}{2\pi\Delta y}\Big)^{K+1}$$

where $\hat{f}(y) = \int_{-\infty}^{\infty} f(t)e(yt)dt$.

We also need to study

$$S_\alpha^h(y) := \sum_{k\le y}\mathfrak{S}(k)k^\alpha \cos(h\log\frac{kx}{y}) - \int_0^y u^\alpha\cos(h\log\frac{ux}{y})du \text{ for } \alpha \ge 0,$$

and

$$T_\alpha^h(y) := \sum_{k>y}\frac{\mathfrak{S}(k)}{k^\alpha}\cos(h\log\frac{kx}{y}) - \int_y^\infty \frac{1}{u^\alpha}\cos(h\log\frac{ux}{y})du \text{ for } \alpha > 1.$$

Then from [4],

$$S_0(y) := S_0^0(y) = -\frac{1}{2}\log y + O((\log y)^{2/3}) = -\frac{1}{2}\log y + \epsilon(y). \tag{3}$$

By partial summation and Lemma 2.7,

$$\begin{aligned} S_\alpha^h(y) =& \epsilon(y)y^\alpha\cos(h\log x) - \frac{\alpha\cos(h\log x)}{2(\alpha^2+h^2)}y^\alpha - \frac{h\sin(h\log x)}{2(\alpha^2+h^2)}y^\alpha \\ &- \int_0^y \epsilon(u)u^{\alpha-1}\Big[\alpha\cos(h\log\frac{ux}{y}) - h\sin(h\log\frac{ux}{y})\Big]du, \end{aligned} \tag{4}$$

and

$$\begin{aligned} T_\alpha^h(y) =& -\frac{\epsilon(y)}{y^\alpha}\cos(h\log x) - \frac{\alpha\cos(h\log x)}{2(\alpha^2+h^2)}\frac{1}{y^\alpha} + \frac{h\sin(h\log x)}{2(\alpha^2+h^2)}\frac{1}{y^\alpha} \\ &+ \int_y^\infty \frac{\epsilon(u)}{u^{\alpha+1}}\Big[\alpha\cos(h\log\frac{ux}{y}) + h\sin(h\log\frac{ux}{y})\Big]du. \end{aligned} \tag{5}$$

Let

$$f(y) := \int_0^y \epsilon(u) - \frac{B}{2}\, du$$

where $B = -C_0 - \log 2\pi$ and C_0 is Euler's constant. Note that

$$f(y) \ll y^{1/2+\epsilon} \tag{6}$$

(see Lemma 2.2 of [3]). From (4) and (5),

$$\begin{aligned} &S_2^h(y)\frac{1}{y^3} + T_2^h(y)y \\ &= -\frac{2\cos(h\log x)}{4+h^2}\frac{1}{y} - \frac{1}{y^3}\int_0^y u\epsilon(u)\Big[2\cos(h\log\frac{ux}{y}) - h\sin(h\log\frac{ux}{y})\Big]du \\ &\quad + y\int_y^\infty \frac{\epsilon(u)}{u^3}\Big[2\cos(h\log\frac{ux}{y}) + h\sin(h\log\frac{ux}{y})\Big]du. \end{aligned} \tag{7}$$

Lemma 4.1.

$$\begin{aligned} I + J =& -\frac{1}{y^3}\int_0^y u\epsilon(u)\Big[2\cos(h\log\frac{ux}{y}) - h\sin(h\log\frac{ux}{y})\Big]du \\ &+ y\int_y^\infty \frac{\epsilon(u)}{u^3}\Big[2\cos(h\log\frac{ux}{y}) + h\sin(h\log\frac{ux}{y})\Big]du \\ =& -\frac{4f(y)}{y}\cos(h\log x) \\ &+ \frac{1}{y^3}\int_0^y f(u)\Big[(2-h^2)\cos(h\log\frac{ux}{y}) - 3h\sin(h\log\frac{ux}{y})\Big]du \\ &+ y\int_y^\infty \frac{f(u)}{u^4}\Big[(6-h^2)\cos(h\log\frac{ux}{y}) + 5h\sin(h\log\frac{ux}{y})\Big]du. \end{aligned}$$

Proof: I can be rewritten as

$$\begin{aligned} &-\frac{1}{y^3}\int_0^y u\Big(\epsilon(u) - \frac{B}{2}\Big)\Big[2\cos(h\log\frac{ux}{y}) - h\sin(h\log\frac{ux}{y})\Big]du \\ &-\frac{B}{2}\frac{1}{y^3}\int_0^y u\Big[2\cos(h\log\frac{ux}{y}) - h\sin(h\log\frac{ux}{y})\Big]du = -I_1 - I_2. \end{aligned}$$

By a substitution $v = \log\frac{ux}{y}$ and Lemma 2.7,

$$I_2 = \frac{B}{2}\frac{1}{y}\cos(h\log x). \tag{8}$$

By integration by parts and (6),

$$\begin{aligned} I_1 =& \frac{f(y)}{y^2}[2\cos(h\log x) - h\sin(h\log x)] \\ &- \frac{1}{y^3}\int_0^y f(u)\Big[(2-h^2)\cos(h\log\frac{ux}{y}) - 3h\sin(h\log\frac{ux}{y})\Big]du. \end{aligned} \tag{9}$$

Similarly, J can be rewritten as

$$\begin{aligned} & y\int_y^\infty \frac{\epsilon(u)-\frac{B}{2}}{u^3}\Big[2\cos{(h\log\frac{ux}{y})}+h\sin{(h\log\frac{ux}{y})}\Big]du \\ +&\frac{B}{2}y\int_y^\infty \frac{1}{u^3}\Big[2\cos{(h\log\frac{ux}{y})}+h\sin{(h\log\frac{ux}{y})}\Big]du = J_1+J_2 \end{aligned}$$

By a substitution $v=\log\dfrac{ux}{y}$ and Lemma 2.7,

$$J_2=\frac{B}{2}\frac{1}{y}\cos{(h\log x)}. \tag{10}$$

By integration by parts and (6),

$$\begin{aligned} J_1 =& -\frac{f(y)}{y^2}[2\cos{(h\log x)}+h\sin{(h\log x)}] \\ &+y\int_y^\infty \frac{f(u)}{u^4}\Big[(6-h^2)\cos{(h\log\frac{ux}{y})}+5h\sin{(h\log\frac{ux}{y})}\Big]du. \end{aligned} \tag{11}$$

(8), (9), (10) and (11) together gives the lemma.

Lemma 4.2.

$$S_2^h(y)\frac{1}{y^3}+T_2^h(y)y=-\frac{2\cos{(h\log x)}}{4+h^2}\frac{1}{y}-\frac{4f(y)}{y^2}\cos{(h\log x)}+G_1(y)+G_2(y)$$

where

$$G_1(y)=\frac{1}{y^3}\int_0^y f(u)\Big[(2-h^2)\cos{(h\log\frac{ux}{y})}-3h\sin{(h\log\frac{ux}{y})}\Big]du,$$

and

$$G_2(y)=y\int_y^\infty \frac{f(u)}{u^4}\Big[(6-h^2)\cos{(h\log\frac{ux}{y})}+5h\sin{(h\log\frac{ux}{y})}\Big]du.$$

Proof: Combine (7) and Lemma 4.1.

Lemma 4.3. *For any integer $n\geq 1$,*

$$\int_1^\infty \frac{1}{y^n}Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)dy=\int_1^\infty \frac{1}{y^n}\frac{\sin\frac{T}{x}y}{\frac{T}{x}y}dy+O\Big(\Delta\log\frac{1}{\Delta}\Big).$$

When $n\neq 2$, the error term can be replaced by $O(\Delta)$.

Proof: This is Lemma 3.3 in [3].

Lemma 4.4. *If $F(y)\ll y^{-3/2+\epsilon}$ for $y\geq 1$, then*

$$\int_1^\infty F(y)Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)dy=\int_1^\infty F(y)\frac{\sin\frac{T}{x}y}{\frac{T}{x}y}dy+O(\Delta).$$

Proof: This is Lemma 3.4 in [3].

5 Proof of Theorem 1.2

Throughout this section, we assume $\tau = T^{1-\epsilon} \leq T/\log^M T \leq x \leq T^{2-2\epsilon}$, $U = \log^M T$ for $M > 2$, $H^* = \tau^{-2} x^{2/(1-\epsilon)}$, and $\Psi_U(t)$ is defined as in the previous section. The implicit constants in the error terms may depend on ϵ, K and M.

Our method is that of Goldston and Gonek [6] and it is very similar to [3]. Let $s = \sigma + it$,

$$A_h(s) := \sum_{n\leq x} \frac{\Lambda(n)(1+n^{ih})}{n^s} \text{ and } A_h^*(s) := \sum_{n>x} \frac{\Lambda(n)(1+n^{ih})}{n^s},$$

$$A(s) := \frac{1}{2}A_0(s) \text{ and } A^*(s) := \frac{1}{2}A_0^*(s).$$

By Lemma 2.4, with slight modifications, one has

$$\begin{aligned} 4\pi F_h(x,T) = &\int_0^T \Big| \frac{1}{x}\Big(A_h(-\frac{1}{2}+it) - \int_1^x (1+u^{ih}) u^{1/2-it} du \Big) \\ &+ x\Big(A_h^*(\frac{3}{2}+it) - \int_x^\infty (1+u^{ih}) u^{-3/2-it} du \Big)\Big|^2 dt \\ -2 &\int_0^T \Big| \frac{1}{x}\Big(A(-\frac{1}{2}+it) - \int_1^x u^{1/2-it} du \Big) \\ &+ x\Big(A^*(\frac{3}{2}+it) - \int_x^\infty u^{-3/2-it} du \Big)\Big|^2 dt + O\big(\tilde{h}\log^3 T\big). \end{aligned}$$

Inserting $\Psi_U(t/T)$ into the integral and extending the range of integration to the whole real line, we have

$$\begin{aligned} 4\pi F(x,T) = &\frac{1}{x^2} I_1(x,T) + x^2 I_2(x,T) - \frac{2}{x^2} I_3(x,T) - 2x^2 I_4(x,T) \\ &+ O\Big(\frac{T(\log T)^2}{U}\Big) + O\Big(\frac{x^{1+6\epsilon}}{T}\Big) \end{aligned} \tag{12}$$

where

$$\begin{aligned} I_1(x,T) &= \int_{-\infty}^{\infty} \Big| A_h(-\frac{1}{2}+it) - \int_1^x (1+u^{ih}) u^{1/2-it} du \Big|^2 \Psi_U\Big(\frac{t}{T}\Big) dt, \\ I_2(x,T) &= \int_{-\infty}^{\infty} \Big| A_h^*(\frac{3}{2}+it) - \int_x^\infty (1+u^{ih}) u^{-3/2-it} du \Big|^2 \Psi_U\Big(\frac{t}{T}\Big) dt, \\ I_3(x,T) &= \int_{-\infty}^{\infty} \Big| A(-\frac{1}{2}+it) - \int_1^x u^{1/2-it} du \Big|^2 \Psi_U\Big(\frac{t}{T}\Big) dt, \\ I_4(x,T) &= \int_{-\infty}^{\infty} \Big| A^*(\frac{3}{2}+it) - \int_x^\infty u^{-3/2-it} du \Big|^2 \Psi_U\Big(\frac{t}{T}\Big) dt \end{aligned}$$

by Lemma 1 of [7] with modification $V = -T/U$ and $T - T/U$, and $W = 2T/U$. The contribution from the cross terms are estimated via Theorem 3 of [6]. Note that by partial summation with the Riemann Hypothesis and **TPC**,

$$\begin{aligned}&\sum_{n\le x}\Lambda(n)(1+n^{ih})=\int_1^x(1+u^{ih})\,du+O(\tilde{h}x^{1/2+\epsilon}),\\&\sum_{n\le x}\Lambda(n)\Lambda(n+k)(1+n^{ih})(1+(n+k)^{-ih})\\=&\mathfrak{S}(k)\int_1^x(1+u^{ih})(1+(u+k))^{-ih}\,du+O(\tilde{h}x^{1/2+\epsilon}).\end{aligned}$$

By Corollary 1 of [6] (see also the calculations at the end of [6] and [7]),

$$\begin{aligned}I_1(x,T)=&\hat{\Psi}_U(0)T\sum_{n\le x}\Lambda^2(n)n|1+n^{ih}|^2\\&+4\pi\Big(\frac{T}{2\pi}\Big)^3\int_{T/2\pi x}^{\infty}\Big[\sum_{k\le 2\pi xv/T}\mathfrak{S}(k)k^2\big(1+(\frac{kT}{2\pi v})^{ih}\big)\\&\times\big(1+(\frac{kT}{2\pi v}+k)^{-ih}\big)\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3}\\&-4\pi\Big(\frac{T}{2\pi}\Big)^3\int_{T/2\pi\tau x}^{\infty}\Big[\int_0^{2\pi xv/T}u^2\big|1+(\frac{uT}{2\pi v})^{ih}\big|^2du\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3}\\&+O\Big(\tilde{h}\frac{x^{3+6\epsilon}}{T}\Big)+O(\tilde{h}x^{5/2+7\epsilon}).\end{aligned}$$

Note that

$$\begin{aligned}&\Big(1+(\frac{kT}{2\pi v})^{ih}\Big)\Big(1+(\frac{kT}{2\pi v}+k)^{-ih}\Big)\\=&\Big|1+(\frac{kT}{2\pi v})^{ih}\Big|^2+\Big(1+(\frac{kT}{2\pi v})^{ih}\Big)\Big((\frac{kT}{2\pi v}+k)^{-ih}-(\frac{kT}{2\pi v})^{-ih}\Big)\\=&\Big|1+(\frac{kT}{2\pi v})^{ih}\Big|^2+\Big(1+(\frac{kT}{2\pi v})^{ih}\Big)\Big(\frac{kT}{2\pi v}\Big)^{-ih}\Big((1+\frac{2\pi v}{T})^{-ih}-1\Big)\\=&\Big|1+(\frac{kT}{2\pi v})^{ih}\Big|^2+O\Big(\min(\frac{hv}{T},1)\Big).\end{aligned}\tag{13}$$

Thus,

$$\begin{aligned}&\int_{T/2\pi x}^{\infty}\Big[\sum_{k\le 2\pi xv/T}\mathfrak{S}(k)k^2\big(1+(\frac{kT}{2\pi v})^{ih}\big)\big(1+(\frac{kT}{2\pi v}+k)^{-ih}\big)\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3}\\=&\int_{T/2\pi x}^{\infty}\Big[\sum_{k\le 2\pi xv/T}\mathfrak{S}(k)k^2\big|1+(\frac{kT}{2\pi v})^{ih}\big|^2\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3}\\&+O\Big(\int_{T/2\pi x}^{T^{2-\epsilon}/x}(\frac{xv}{T})^3\frac{hv}{T}\frac{1}{v}\frac{dv}{v^3}+\int_{T^{2-\epsilon}/x}^{\infty}(\frac{xv}{T})^3\frac{1}{\Delta v^2}\frac{dv}{v^3}\Big)\\=&\int_{T/2\pi x}^{\infty}\Big[\sum_{k\le 2\pi xv/T}\mathfrak{S}(k)k^2\big|1+(\frac{kT}{2\pi v})^{ih}\big|^2\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3}+O\Big(\frac{hx^2}{T^{2+\epsilon}}+\frac{x^4}{\Delta T^{5-\epsilon}}\Big)\end{aligned}$$

as $\sum\limits_{k\leq x}\mathfrak{S}(k)\sim x$ and $Re\hat{\Psi}_U(v)\ll \min\,(\frac{1}{v},\frac{1}{\Delta v^2})$. Therefore,

$$\begin{aligned}
I_1(x,T) =& T\sum_{n\leq x}\Lambda^2(n)n(2+2\cos{(h\log n)}) \\
&+4\pi\Big(\frac{T}{2\pi}\Big)^3\int_{T/2\pi x}^{\infty}\Big[\sum_{k\leq 2\pi xv/T}\mathfrak{S}(k)k^2\big(2+2\cos{(h\log\frac{kT}{2\pi v}})\big) \\
&-\int_0^{2\pi xv/T}u^2\big(2+2\cos{(h\log\frac{uT}{2\pi v}})\big)du\Big]Re\hat{\Psi}_U(v)\frac{dv}{v^3} \\
&-4\pi\Big(\frac{T}{2\pi}\Big)^3\int_0^{T/2\pi x}\int_0^{2\pi xv/T}u^2\big(2+2\cos{(h\log\frac{uT}{2\pi v}})\big)duRe\hat{\Psi}_U(v)\frac{dv}{v^3} \\
&+O\Big(\frac{\tilde{h}x^{3+6\epsilon}}{T}\Big)+O(\tilde{h}x^{5/2+7\epsilon})+O(\tilde{h}x^2T^{1-\epsilon})+O\Big(\frac{x^4}{\Delta T^{2-\epsilon}}\Big).
\end{aligned}$$

Similarly, by Corollary 2 of [6],

$$\begin{aligned}
I_2(x,T) =& T\sum_{x<n}\frac{\Lambda^2(n)}{n^3}(2+2\cos{(h\log n)}) \\
&+\frac{8\pi^2}{T}\int_0^{TH^*/2\pi x}\Big[\sum_{2\pi xv/T\leq k\leq H^*}\frac{\mathfrak{S}(k)}{k^2}\big(2+2\cos{(h\log\frac{kT}{2\pi v}})\big) \\
&-\int_{2\pi xv/T}^{H^*}\frac{1}{u^2}\big(2+2\cos{(h\log\frac{uT}{2\pi v}})\big)du\Big]Re\hat{\Psi}_U(v)vdv \\
&+O(\tilde{h}T^{-1}x^{-1+6\epsilon})+O(\tilde{h}x^{-3/2+7\epsilon})+O(\tilde{h}T^{1-\epsilon/2}x^{-2})+O\Big(\frac{\tilde{h}H^*}{\Delta x^2}\Big)
\end{aligned}$$

where the last error term comes from the error term in (13). $I_3(x,T)$ and $I_4(x,T)$ are computed in [3] or one can simply set $h=0$ in $I_1(x,T)$ and $I_2(x,T)$, and divide by 4. Putting these into (12) with a substitution $y=\dfrac{2\pi xv}{T}$ and using

Lemma 2.8,

$$
\begin{aligned}
&4\pi F_h(x,T)\\
&= 2T\Big[\frac{4\cos(h\log x)}{4+h^2}\log x - \frac{8h\sin(h\log x)}{(4+h^2)^2}\Big]\\
&+ 4T\int_1^\infty \Big[\sum_{k\le y}\mathfrak{S}(k)k^2\cos(h\log\frac{kx}{y}) - \int_0^y u^2\cos(h\log\frac{ux}{y})du\Big] Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)\frac{dy}{y^3}\\
&- 4T\int_0^1\int_0^y u^2\cos(h\log\frac{ux}{y})du Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)\frac{dy}{y^3}\\
&+ 4T\int_1^{H^*}\Big[\sum_{y\le k\le H^*}\frac{\mathfrak{S}(k)\cos(h\log\frac{kx}{y})}{k^2} - \int_y^{H^*}\frac{\cos(h\log\frac{ux}{y})}{u^2}du\Big] Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)ydy\\
&+ 4T\int_0^1\Big[\sum_{k\le H^*}\frac{\mathfrak{S}(k)\cos(h\log\frac{kx}{y})}{k^2} - \int_y^{H^*}\frac{\cos(h\log\frac{ux}{y})}{u^2}du\Big] Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)ydy\\
&+ O\Big(\frac{\tilde{h}x^{1+6\epsilon}}{T}\Big) + O(\tilde{h}x^{1/2+7\epsilon}) + O\Big(\frac{\tilde{h}x^2}{T^{2-2\epsilon}}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2}T}\Big)
\end{aligned}
$$

From Lemma 2.7,

$$
\int e^{ax}\cos bx = \frac{a}{a^2+b^2}e^{ax}\cos bx + \frac{b}{a^2+b^2}e^{ax}\sin bx. \tag{14}
$$

Also,

$$
\begin{aligned}
\int_0^1 Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)dy =& \frac{x}{T}\int_0^{T/x}\frac{\sin u}{u}(1+O(\Delta^2u^2))du\\
=& \frac{x}{T}\int_0^{T/x}\frac{\sin u}{u}du + O\Big(\frac{\Delta^2 T}{x}\Big).
\end{aligned} \tag{15}
$$

By appropriate change of variables, (14) and (15),

$$
\begin{aligned}
&\int_0^1\int_0^y u^2\cos(h\log\frac{ux}{y})du Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)\frac{dy}{y^3}\\
=&\frac{x}{T}\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{3}{9+h^2}\cos(h\log x) + \frac{h}{9+h^2}\sin(h\log x)\Big] + O\Big(\frac{\Delta^2 T}{x}\Big).\\
&\int_0^1\Big[\sum_{k\le H^*}\frac{\mathfrak{S}(k)\cos(h\log\frac{kx}{y})}{k^2} - \int_y^{H^*}\frac{\cos(h\log\frac{ux}{y})}{u^2}du\Big] Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)ydy\\
=&\sum_{k=1}^\infty\frac{\mathfrak{S}(k)}{k^2}\int_0^1 y\cos(h\log\frac{kx}{y})\frac{\sin\frac{Ty}{x}}{\frac{Ty}{x}}dy + O\Big(\frac{1}{H^*}\Big) + O\Big(\frac{\Delta^2 T}{x}\Big)\\
&-\frac{x}{T}\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{1}{1+h^2}\cos(h\log x) - \frac{h}{1+h^2}\sin(h\log x)\Big] + O\Big(\frac{\Delta^2 T}{x}\Big).
\end{aligned}
$$

Therefore, with the notation $S^h_\alpha(y)$ and $T^h_\alpha(y)$,

$$\begin{aligned}
&4\pi F_h(x,T)\\
&= 2T\Big[\frac{4\cos{(h\log x)}}{4+h^2}\log x - \frac{8h\sin{(h\log x)}}{(4+h^2)^2}\Big]\\
&+ 4T\int_1^\infty S^h_2(y) Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big)\frac{dy}{y^3}\\
&+ 4T\int_1^{H^*} \big(T^h_2(y) - T^h_2(H^*)\big) Re\hat{\Psi}_U\Big(\frac{Ty}{2\pi x}\Big) y dy\\
&- 4x\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{3\cos{(h\log x)}}{9+h^2} + \frac{h\sin{(h\log x)}}{9+h^2}\Big]\\
&- 4x\int_0^{T/x}\frac{\sin u}{u}du\Big[\frac{\cos{(h\log x)}}{1+h^2} - \frac{h\sin{(h\log x)}}{1+h^2}\Big]\\
&+ 4T\sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2}\int_0^1 y\cos{(h\log\frac{kx}{y})}\frac{\sin\frac{Ty}{x}}{\frac{Ty}{x}}dy\\
&+ O\Big(\frac{\tilde{h}x^{1+6\epsilon}}{T}\Big) + O(\tilde{h}x^{1/2+7\epsilon}) + O\Big(\frac{\tilde{h}x^2}{T^{2-2\epsilon}}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2}T}\Big).
\end{aligned}$$

By (3) and (5), $T^h_2(H^*) \ll \dfrac{h(\log H^*)^{2/3}}{(H^*)^2}$. It follows that the contribution from $T^h_2(H^*)$ in the second integral is $O(hT^{-\epsilon})$. Also, one can extend the upper limit of the second integral to ∞ with an error $O(hT^{-\epsilon})$ by (3) and (5) again. Finally, we obtain the theorem by applying Lemma 4.2, Lemma 4.3, Lemma 4.4, (6) and dividing by 4π.

6 Proof of Theorem 1.3 and 1.4

Proof of Theorem 1.3: It follows directly from Theorem 1.2 by observing that all the other main terms besides the first one are $O(x)$ because of (6).

Before proving Theorem 1.4, we need the following lemmas.

Lemma 6.1.

$$\int_1^\infty \frac{\sin ax}{x^{2n}}dx = \frac{a^{2n-1}}{(2n-1)!}\Big[\sum_{k=1}^{2n-1}\frac{(2n-k-1)!}{a^{2n-k}}\sin\big(a+(k-1)\frac{\pi}{2}\big) + (-1)^n ci(a)\Big]$$

where $ci(x) = -\int_x^\infty \frac{\cos t}{t}dt = C_0 + \log x + \int_0^x \frac{\cos t - 1}{t}dt$ *and* C_0 *is Euler's constant.*

Proof: This is formula 3.761(3) on P.430 of [8] which can be proved by integration by parts repeatedly.

Lemma 6.2. *If* $F(y) \ll y^{-3/2+\epsilon}$ *for* $y \geq 1$*, then for* $T \leq x$*,*

$$\int_1^\infty F(y) \frac{\sin \frac{T}{x} y}{\frac{T}{x} y} dy = \int_1^\infty F(y) dy + O\Big((\frac{T}{x})^{1/2-\epsilon} \Big).$$

Proof: This is Lemma 5.2 in [3].

Lemma 6.3.

$$\begin{aligned} I &= \int_1^\infty \frac{1}{y^3} \int_0^y f(u) \Big[(2-h^2) \cos (h \log \frac{ux}{y}) - 3h \sin (h \log \frac{ux}{y}) \Big] du\, dy \\ &= \int_0^1 f(u) [\cos (h \log ux) - h \sin (h \log ux)]\, du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2}\, du [\cos (h \log x) - h \sin (h \log x)]. \end{aligned}$$

Proof: Because of (6), we can change the order of integration.

$$\begin{aligned} I &= \int_0^1 f(u) \int_1^\infty \frac{1}{y^3} \Big[(2-h^2) \cos (h \log \frac{ux}{y}) - 3h \sin (h \log \frac{ux}{y}) \Big] dy\, du \\ &\quad + \int_1^\infty f(u) \int_u^\infty \frac{1}{y^3} \Big[(2-h^2) \cos (h \log \frac{ux}{y}) - 3h \sin (h \log \frac{ux}{y}) \Big] dy\, du \\ &= \int_0^1 f(u) \Big\{ (2-h^2) \Big[\frac{2}{4+h^2} \cos (h \log ux) + \frac{h}{4+h^2} \sin (h \log ux) \Big] \\ &\quad - 3h \Big[\frac{2}{4+h^2} \sin (h \log ux) - \frac{h}{4+h^2} \cos (h \log ux) \Big] \Big\} du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2} \Big\{ (2-h^2) \Big[\frac{2}{4+h^2} \cos (h \log x) + \frac{h}{4+h^2} \sin (h \log x) \Big] \\ &\quad - 3h \Big[\frac{2}{4+h^2} \sin (h \log x) - \frac{h}{4+h^2} \cos (h \log x) \Big] \Big\} du. \end{aligned}$$

by substituting $v = \log \frac{ux}{y}$ and applying Lemma 2.7. The lemma follows after some simple algebra.

Lemma 6.4.

$$\begin{aligned} J &= \int_1^\infty y \int_y^\infty \frac{f(u)}{u^4} \Big[(6-h^2) \cos (h \log \frac{ux}{y}) + 5h \sin (h \log \frac{ux}{y}) \Big] du\, dy \\ &= - \int_1^\infty \frac{f(u)}{u^4} [3 \cos (h \log ux) + h \sin (h \log ux)]\, du \\ &\quad + \int_1^\infty \frac{f(u)}{u^2}\, du [3 \cos (h \log x) + h \sin (h \log x)]. \end{aligned}$$

Proof: Again, because of (6), we can change the order of integration.

$$
\begin{aligned}
J =& \int_1^\infty \int_1^u y\Big[(6-h^2)\cos(h\log\frac{ux}{y}) + 5h\sin(h\log\frac{ux}{y})\Big]\,dy\,du \\
=& \int_1^\infty \frac{f(u)}{u^4}\Big\{(6-h^2)\Big[\frac{-2}{4+h^2}\cos(h\log ux) + \frac{h}{4+h^2}\sin(h\log ux)\Big] \\
&+ 5h\Big[\frac{-2}{4+h^2}\sin(h\log ux) - \frac{h}{4+h^2}\cos(h\log ux)\Big] \\
&- (6-h^2)\Big[\frac{-2}{4+h^2}\cos(h\log x) + \frac{h}{4+h^2}\sin(h\log x)\Big] \\
&- 5h\Big[\frac{-2}{4+h^2}\sin(h\log x) - \frac{h}{4+h^2}\cos(h\log x)\Big]\Big\}du.
\end{aligned}
$$

by substituting $v = \log\frac{ux}{y}$ and applying Lemma 2.7. The lemma follows after some simple algebra.

Lemma 6.5.

$$
\begin{aligned}
S =& \sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2}\int_0^1 y\cos(h\log\frac{kx}{y})\,dy \\
=& \Big[\frac{1}{1+h^2}\cos(h\log x) - \frac{h}{1+h^2}\sin(h\log x)\Big] \\
&- \Big[\frac{4-h^2}{2(4+h^2)^2}\cos(h\log x) - \frac{2h}{(4+h^2)^2}\sin(h\log x)\Big] \\
&+ \frac{B}{2}\Big[\frac{2}{4+h^2}\cos(h\log x) - \frac{h}{4+h^2}\sin(h\log x)\Big] + \Big(1+\frac{B}{2}\Big)\cos(h\log x) \\
&+ \int_1^\infty \frac{f(u)}{u^4}\Big[3\cos(h\log ux) + h\sin(h\log ux)\Big]du.
\end{aligned}
$$

Proof: By substituting $v = \log\frac{kx}{y}$ and Lemma 2.7,

$$
S = \frac{2}{4+h^2}\sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2}\cos(h\log kx) - \frac{h}{4+h^2}\sum_{k=1}^\infty \frac{\mathfrak{S}(k)}{k^2}\sin(h\log kx).
$$

Recall the definition of $S_0(u)$ from (3) and use partial summation,

$$\begin{aligned}
S =&\frac{2}{4+h^2}\int_1^\infty \frac{S_0(u)+u}{u^3}[2\cos(h\log ux)+h\sin(h\log ux)]\,du\\
&-\frac{h}{4+h^2}\int_1^\infty \frac{S_0(u)+u}{u^3}[-h\cos(h\log ux)+2\sin(h\log ux)]\,du\\
=&\int_1^\infty \frac{S_0(u)+u}{u^3}\cos(h\log ux)\,du\\
=&\int_1^\infty \frac{u-\frac{1}{2}\log u+\epsilon(u)}{u^3}\cos(h\log ux)\,du\\
=&\int_1^\infty \frac{1}{u^2}\cos(h\log ux)\,du-\frac{1}{2}\int_1^\infty \frac{\log u}{u^3}\cos(h\log ux)\,du\\
&+\frac{B}{2}\int_1^\infty \frac{1}{u^3}\cos(h\log ux)\,du+\int_1^\infty \frac{\epsilon(u)-\frac{B}{2}}{u^3}\cos(h\log ux)\,du\\
=&I_1-\frac{1}{2}I_2+\frac{B}{2}I_3+I_4.
\end{aligned}$$

By appropriate substitution and Lemma 2.7,

$$\begin{aligned}
I_1 =&\frac{1}{1+h^2}\cos(h\log x)-\frac{h}{1+h^2}\sin(h\log x),\\
I_2 =&\frac{4-h^2}{(4+h^2)^2}\cos(h\log x)-\frac{2h}{(4+h^2)^2}\sin(h\log x),\\
I_3 =&\frac{2}{4+h^2}\cos(h\log x)-\frac{h}{4+h^2}\sin(h\log x).
\end{aligned}$$

Finally, by integration by parts,

$$\begin{aligned}
I_4 =&\int_1^\infty \frac{\cos(h\log ux)}{u^3}df(u)\\
=&\Big(1+\frac{B}{2}\Big)\cos(h\log x)+\int_1^\infty \frac{f(u)}{u^4}[3\cos(h\log ux)+h\sin(h\log ux)]\,du
\end{aligned}$$

because $f(1)=-1-\frac{B}{2}$. Combining the results for I_1, I_2, I_3 and I_4, we have the lemma.

Lemma 6.6.

$$\begin{aligned}
&\int_0^1 f(u)[\cos(h\log ux)-h\sin(h\log ux)]\,du\\
=&-\frac{1}{2}\Big[\frac{4+3h^2}{(4+h^2)^2}\cos(h\log x)+\frac{h^3}{(4+h^2)^2}\sin(h\log x)\Big]\\
&-\Big(\frac{1}{2}+\frac{B}{2}\Big)\Big[\frac{2+h^2}{4+h^2}\cos(h\log x)-\frac{h}{4+h^2}\sin(h\log x)\Big]\\
&-\frac{1}{2}\Big[\frac{3+h^2}{9+h^2}\cos(h\log x)-\frac{2h}{9+h^2}\sin(h\log x)\Big].
\end{aligned}$$

Proof: The key is $\epsilon(u) = \frac{1}{2}\log u - u$ when $0 \le u \le 1$ (see (3)). So,

$$f(u) = \int_0^u \epsilon(v) - \frac{B}{2}\, dv = \frac{1}{2} u \log u - \Big(\frac{1}{2} + \frac{B}{2}\Big) u - \frac{1}{2} u^2.$$

Putting this into the integral and evaluating the integral piece by piece with suitable substitution and Lemma 2.7, one gets the lemma.

Proof of Theorem 1.4: First observe that when $T \le x \le T^{2-29\epsilon}$, the error terms in Theorem 1.2 is $O(\frac{hT}{\log^{M-2} T})$. Rewrite Theorem 1.2 as

$$F_h(x,T) = T_1 + T_2 + T_3 + T_4 + T_5 + O\Big(\frac{\tilde{h}T}{\log^{M-2} T}\Big).$$

Since $\frac{\sin u}{u} = 1 + O(u^2)$,

$$\begin{aligned} T_3 &= -\frac{T}{\pi}\Big[\frac{3\cos(h\log x)}{9+h^2} + \frac{h\sin(h\log x)}{9+h^2}\Big] + O\Big(T\big(\frac{T}{x}\big)^2\Big), \\ T_4 &= -\frac{T}{\pi}\Big[\frac{\cos(h\log x)}{1+h^2} - \frac{h\sin(h\log x)}{1+h^2}\Big] + O\Big(T\big(\frac{T}{x}\big)^2\Big). \end{aligned}$$

By Lemma 6.5,

$$\begin{aligned} T_5 =& \frac{T}{\pi}\sum_{k=1}^{\infty}\frac{\mathfrak{S}(k)}{k^2}\int_0^1 y\cos\big(h\log\frac{kx}{y}\big)\, dy + O\Big(T\big(\frac{T}{x}\big)^2\Big) \\ =& \frac{T}{\pi}\Big[\frac{\cos(h\log x)}{1+h^2} - \frac{h\sin(h\log x)}{1+h^2}\Big] \\ &- \frac{T}{\pi}\Big[\frac{4-h^2}{2(4+h^2)^2}\cos(h\log x) - \frac{2h}{(4+h^2)^2}\sin(h\log x)\Big] \\ &+ \frac{T}{\pi}\frac{B}{2}\Big[\frac{2\cos(h\log x)}{4+h^2} - \frac{h\sin(h\log x)}{4+h^2}\Big] + \frac{T}{\pi}\Big(1+\frac{B}{2}\Big)\cos(h\log x) \\ &+ \frac{T}{\pi}\int_1^{\infty}\frac{f(u)}{u^4}\Big[3\cos(h\log ux) + h\sin(h\log ux)\Big] du + O\Big(T\big(\frac{T}{x}\big)^2\Big). \end{aligned}$$

By Lemma 4.3, (6), Lemma 6.2, Lemma 6.3, Lemma 6.4 and Lemma 6.6,

$$\begin{aligned}
T_2 =& -\frac{2x\cos(h\log x)}{\pi(4+h^2)}\int_1^\infty \frac{\sin\frac{Ty}{x}}{y^2}dy - \frac{4T}{\pi}\cos(h\log x)\int_1^\infty \frac{f(y)}{y^2}dy \\
&+ \frac{T}{\pi}\int_1^\infty \big(G_1(y)+G_2(y)\big)\,dy + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) + O\Big(\frac{\tilde{h}T}{\log^M T}\Big) \\
=& -\frac{2x\cos(h\log x)}{\pi(4+h^2)}\int_1^\infty \frac{\sin\frac{Ty}{x}}{y^2}dy + O\Big(\frac{\tilde{h}T}{\log^M T}\Big) \\
&+ \frac{T}{\pi}\int_0^1 f(u)[\cos(h\log ux) - h\sin(h\log ux)]\,du \\
&- \frac{T}{\pi}\int_1^\infty \frac{f(u)}{u^4}[3\cos(h\log ux) + h\sin(h\log ux)]\,du + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) \\
=& -\frac{2x\cos(h\log x)}{\pi(4+h^2)}\int_1^\infty \frac{\sin\frac{Ty}{x}}{y^2}dy + O\Big(\frac{\tilde{h}T}{\log^M T}\Big) \\
&- \frac{T}{\pi}\frac{1}{2}\Big[\frac{4+3h^2}{(4+h^2)^2}\cos(h\log x) + \frac{h^3}{(4+h^2)^2}\sin(h\log x)\Big] \\
&- \frac{T}{\pi}\Big(\frac{1}{2}+\frac{B}{2}\Big)\Big[\frac{2+h^2}{4+h^2}\cos(h\log x) - \frac{h}{4+h^2}\sin(h\log x)\Big] \\
&- \frac{T}{\pi}\frac{1}{2}\Big[\frac{3+h^2}{9+h^2}\cos(h\log x) - \frac{2h}{9+h^2}\sin(h\log x)\Big] \\
&- \frac{T}{\pi}\int_1^\infty \frac{f(u)}{u^4}[3\cos(h\log ux) + h\sin(h\log ux)]\,du + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big).
\end{aligned}$$

Therefore, with miraculous cancellations,

$$\begin{aligned}
T_2+T_3+T_4+T_5 =& -\frac{2x\cos(h\log x)}{\pi(4+h^2)}\int_1^\infty \frac{\sin\frac{Ty}{x}}{y^2}dy + \frac{T}{\pi}\frac{2B\cos(h\log x)}{4+h^2} \\
&+ \frac{T}{\pi}\frac{4h\sin(h\log x)}{(4+h^2)^2} + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) + O\Big(\frac{\tilde{h}T}{\log^M T}\Big).
\end{aligned}$$

By Lemma 6.1 and $B = -C_0 - \log 2\pi$,

$$\begin{aligned}
F_h(x,T) =& \frac{T}{\pi}\Big[\frac{2\cos(h\log x)}{4+h^2}\log x\Big] - \frac{2T\cos(h\log x)}{\pi(4+h^2)}\Big[\frac{\sin(T/x)}{T/x} - ci\big(\frac{T}{x}\big)\Big] \\
&+ \frac{T}{\pi}\frac{2B\cos(h\log x)}{4+h^2} + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2} T}\Big) \\
=& \frac{T}{\pi}\Big[\frac{2\cos(h\log x)}{4+h^2}\log x\Big] - \frac{2T\cos(h\log x)}{\pi(4+h^2)}\Big[1 - C_0 - \log\frac{T}{x} \\
&+ C_0 + \log 2\pi\Big] + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2} T}\Big) \\
=& \frac{T}{2\pi}\log\frac{T}{2\pi e}\Big[\frac{4\cos(h\log x)}{4+h^2}\Big] + O\Big(T\big(\frac{T}{x}\big)^{1/2-\epsilon}\Big) + O\Big(\frac{\tilde{h}T}{\log^{M-2} T}\Big).
\end{aligned}$$

7 Sketch for Conjecture 1.2

Fix $\alpha > 0$. Let $r(u)$ be an even function which is almost the characteristic function of the interval $[-\alpha, \alpha]$ with $\hat{r}(\alpha) \ll \frac{1}{\alpha^2}$ (see page 87 of [1] for detail construction). We use Conjecture 1.1 to compute the right hand side of (1).

$$
\begin{aligned}
I =& \int_{-\infty}^{\infty} F_h(\alpha)\hat{r}(\alpha)d\alpha = 2\int_{0}^{\infty} F_h(\alpha)\hat{r}(\alpha)d\alpha \\
=& 2(1+o(1))\log T \int_0^1 T^{-2\alpha}\hat{r}(\alpha)d\alpha + 2\frac{4}{4+h^2}\int_0^1 \alpha\cos(h\log T\alpha)\hat{r}(\alpha)d\alpha \\
&+ 2\frac{4}{4+h^2}\int_1^{\infty}\cos(h\log T\alpha)\hat{r}(\alpha)d\alpha + O\Big(\frac{1}{A}\Big) + o(1) \\
=& \frac{4}{4+h^2}\int_{-\infty}^{\infty}\cos(h\log T\alpha)\hat{r}(\alpha)d\alpha - \frac{4}{4+h^2}\int_{-1}^{1}(1-|\alpha|)\cos(h\log T\alpha)\hat{r}(\alpha)d\alpha \\
&+ (1+o(1))\log T\int_{-\infty}^{\infty} T^{-2|\alpha|}\hat{r}(\alpha)d\alpha + O\Big(\frac{1}{A}\Big) + o(1) \\
=& \frac{4}{4+h^2}\int_{-\infty}^{\infty}\hat{r}_1(\alpha)d\alpha - \frac{4}{4+h^2}\int_{-1}^{1}(1-|\alpha|)\hat{r}_1(\alpha)d\alpha \\
&+ (1+o(1))\log T\int_{-\infty}^{\infty} T^{-2|\alpha|}\hat{r}(\alpha)d\alpha + O\Big(\frac{1}{A}\Big) + o(1) \\
=& \frac{4}{4+h^2}I_1 - \frac{4}{4+h^2}I_2 + (1+o(1))I_3 + O\Big(\frac{1}{A}\Big) + o(1)
\end{aligned}
$$

where $r_1(u) = r(u + \frac{h\log T}{2\pi})$. As $\int_{-\infty}^{\infty}\hat{r}_1(\alpha)d\alpha = r_1(0)$,

$$
I_1 = r_1(0) = r\Big(\frac{h\log T}{2\pi}\Big).
$$

By $\int f\hat{g} = \int \hat{f}g$, the transform pair and the definition of $r(u)$,

$$
f(t) = \max(1-|t|, 0), \quad \hat{f}(u) = \Big(\frac{\sin\pi u}{\pi u}\Big)^2,
$$

$$
I_2 = \int_{-\infty}^{\infty} r_1(u)\Big(\frac{\sin\pi u}{\pi u}\Big)^2 du = \int_{-\alpha+h\log T/(2\pi)}^{\alpha+h\log T/(2\pi)}\Big(\frac{\sin\pi u}{\pi u}\Big)^2 du + o(1).
$$

Similarly, by the transform pair

$$
f(t) = e^{-2a|t|}, \quad \hat{f}(u) = \frac{4a}{4a^2+(2\pi u)^2},
$$

$$
I_3 = \int_{-\alpha}^{\alpha}\frac{4\log^2 T}{4\log^2 T+(2\pi u)^2}du + o(1) = \int_{-\alpha+h\log T/(2\pi)}^{\alpha+h\log T/(2\pi)} 1du + o(1).
$$

Therefore,

$$
I = \frac{4}{4+h^2}r\Big(\frac{h\log T}{2\pi}\Big) + \int_{-\alpha+h\log T/(2\pi)}^{\alpha+h\log T/(2\pi)} 1 - \frac{4}{4+h^2}\Big(\frac{\sin\pi u}{\pi u}\Big)^2 du + O\Big(\frac{1}{A}\Big) + o(1). \tag{16}
$$

Now, the left hand side of (1) is

$$\frac{4}{4+h^2} r\Big(\frac{h \log T}{2\pi}\Big) + \Big(\frac{T}{2\pi}\log T\Big)^{-1} \sum_{\substack{0<\gamma\neq\gamma'\leq T \\ |\gamma-\gamma'-h|\leq 2\pi\alpha/\log T}} (1+o(1)). \tag{17}$$

Combining (16) and (17), we have Conjecture 1.2 by making A arbitrarily large. The only shaky point in the above argument is the error analysis. All of these become rigorous following page $87-90$ of [1].

References

[1] T.H. Chan, *Pair Correlation and Distribution of Prime Numbers*, Thesis, U. of Mich., Ann Arbor, 2002.

[2] T.H. Chan, *On a conjecture of Liu and Ye*, to appear.

[3] T.H. Chan, *More precise Pair Correlation Conjecture on the zeros of the Riemann zeta function*, submitted.

[4] J.B. Friedlander and D.A. Goldston, *Some singular series averages and the distribution of Goldbach numbers in short intervals*, Illinois J. Math. **39** (1995), 158-180.

[5] D.A. Goldston, *Large Differences between Consecutive Prime Numbers*, Thesis, U. of Calif., Berkeley, 1981.

[6] D.A. Goldston and S.M. Gonek, *Mean value theorems for long Dirichlet polynomials and tails of Dirichlet series*, Acta Arith. (2) **84** (1998), 155-192.

[7] D.A. Goldston, S.M. Gonek, A.E. Ozluk and C. Synder, *On the pair correlation of zeros of the Riemann zeta-function*, Proc. London Math. Soc. (3) **80** (2000), 31-49.

[8] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press, sixth edition, 2000.

[9] H.L. Montgomery, *The pair correlation of zeros of the zeta function*, Analytic Number Theory (St. Louis Univ., 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc., Providence, 1973, pp. 181-193.

[10] H.L. Montgomery and R.C. Vaughan, *Hilbert's inequality*, J. London Math. Soc. (2) **8** (1974), 73-82.

Tsz Ho Chan
Case Western Reserve University
Mathematics Department, Yost Hall 220
10900 Euclid Avenue

Cleveland, OH 44106-7058
USA
txc50@po.cwru.edu